



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 9/68, 7/16, 7/18, 7/00, 9/28, 9/127, 9/20, 33/04, 31/355, 31/34, 31/07</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/28977</b> <b>(43) International Publication Date:</b> 25 May 2000 (25.05.00)
<b>(21) International Application Number:</b> PCT/US99/22507 <b>(22) International Filing Date:</b> 29 September 1999 (29.09.99) <b>(30) Priority Data:</b> 09/163,101 29 September 1998 (29.09.98) US <b>(71) Applicant:</b> THIONE INTERNATIONAL, INC. [US/US]; 3201 Andrews Court N.W., Atlanta, GA 30305 (US). <b>(72) Inventor:</b> HERSH, Theodore; 3201 Andrews Court N.W., Atlanta, GA 30305 (US). <b>(74) Agents:</b> WITTENBERG, Malcolm, B. et al.; Crosby, Heafey, Roach & May, Suite 1900, 4 Embarcadero Center, San Francisco, CA 94111-4106 (US).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> ANTIOXIDANT GEL FOR GINGIVAL CONDITIONS  <b>(57) Abstract</b>  A gel, paste, gum or lozenge composition for oral application to human gums to prevent and reduce symptoms of gum disease. The composition includes reduced glutathione and a source of selenium.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

5

**ANTIOXIDANT GEL FOR GINGIVAL CONDITIONS****RELATED APPLICATIONS**

The present application is a continuation-in-part of U.S. Patent application serial no. 08/982,058, filed on December 1, 1997.

**TECHNICAL FIELD OF THE INVENTION**

10

The present invention deals with the combination of several synergistic antioxidants, enzymatic co-factors and amino acids in appropriate delivery vehicles forming gels, pastes, gums or lozenges. The gels, pastes, gums or lozenges are intended to be applied or taken orally to prophylactically prevent gum disease and remedially to a patient to reduce reactive oxygen and other free radical species which are causative inflammatory factors in establishing and promoting such gingival diseases.

15

**BACKGROUND OF THE INVENTION**

Dentists and other professionals involved with the science of oral hygiene have dealt with a number of symptoms or pathologies of the mouth, gum and teeth. Some of these pathologies are initiated and exacerbated through tobacco consumption, whether smoked or chewed, as smokers often suffer from leukoplakia, a white patch on the buccal mucosa. Although leukoplakia is a benign oral lesion, it has a malignant potential requiring a biopsy of the lesion to rule out cancer. Smokers also have more dental tartar (calculus) than non-smokers and often suffer from halitosis.

25

It is known that tooth loss can directly result from untreated destructive periodontal (gum) disease. Further, acute necrotizing ulcerative gingivitis is a destructive, painful inflammatory condition which is, again, more acutely observed in cigarette smokers and tobacco users.

30

Besides leukoplakia, another type of generalized whitish hue on the buccal mucosa is associated with oral submucous fibrosis. This disease occurs

5 mainly in India and Pakistan and their emigres to the western world and is a chronic, progressive, pre-malignant condition, related to tobacco and betel nut chewing.

Periodontal disease, which includes inflammation of the gums, is one of the most prevalent health problems in the world and is the major cause of tooth  
10 loss in the adult population.

Periodontal disease is composed of two entities:

- a) Gingivitis: Inflammation is confined to the gums (gingival tissue).
- b) Periodontitis: The inflammation and infection is present both in the  
15 gingiva and in the connective tissue which supports the teeth.

There is initially an inflammation in the local blood vessel walls (vasculitis) which accounts for the most common symptom of bleeding gums. This is accompanied by migration of white blood cells (leucocytes) to help combat the tissue invasion by pathogenic mouth bacteria. In this inflammatory reaction,  
20 most cells liberate histamine. Inflammatory (white blood cells) also locally generate countless free radicals and anti-enzyme molecules, both of which contribute to tissue pathology. Collagen production may also then be interfered with due to local Vitamin C deficiency, which contributes to the swelling (edema) and redness (erythema) of gingival tissues.

25 Chewing tobacco and the areca nut are associated with higher frequencies of periodontal disease, oral submucous fibrosis and oral malignancies. Betel Quid chewing is common in India, Southeast Asia and the South Pacific Islands. Free radical damage induced by these deleterious alkaloids initiate an inflammatory reaction in the oral cavity. Studies in Taiwan  
30 reveal that arecoline, the major alkaloid in areca nut interferes with the migration, attachment and growth of human gingival fibroblasts. Arecoline also inhibits these cells' ability to synthesize collagen and all these inhibitory effects have been shown to be associated with an intracellular depletion of L-

5 glutathione (GSH). The repeated and long term exposure to Betel Quid's cytotoxic arecoline and tobacco chewing renders these abusers to periodontal and other oral disease, in part due to losses of the locally vital protective antioxidant and detoxificant, GSH.

Prencipe et al. disclose an anti-bacterial, anti-plaque dentifrice in their  
10 U.S. Patent No. 5,424,059. The '059 patent teaches that plaque can be inhibited and that gingivitis and cavities can be reduced or inhibited by applying a dentifrice in the form of a white water soluble alkaline earth metal polishing agent such as dicalcium phosphate or dimagnesium pyrophosphate used together with an anti-bacterial agent and the polyol, xylitol.

15 Galiana Arano in WO96/38122 discloses a novel dentifrice composition in the form of a tablet for buccal hygiene which does not require a toothbrush, toothpaste or dental powder. The tablet comprises an abrasive and foaming agent, sugar, one or more lubricants, a fluoride source and sweeteners. It is taught that the cleansing effect of water is obtained from the saliva stimulated  
20 by sugars while the mechanical effect of "brushing" is obtained from the granules emanating from the destruction of the tablet.

However, prior attempts to treat gum disease have failed for they have not recognized the role played by free radicals. For example, cigarette tar contains high concentrations of free radicals. The most common oxidants  
25 include semiquinone which is in equilibrium with hydroquinones and quinones, particularly in the viscous tar matrix. Many tar extracts are water soluble and reduce oxygen to the superoxide radical which can dismutate to form  $H_2O_2$ . Importantly, glass-fiber type cigarette filters retain almost all of the tar particles that are larger than 0.1 micron. Thus, the filter acts as a trap for tars and  
30 cigarette smoke. There are an inordinately large number of free radicals, greater than  $10^{15}$ , in each puff of the gas phase of cigarette smoke. While the oxidants in tar are stable, those organic radicals in the gas phase smoke are reactive carbon and oxygen centered radicals with extremely short half-lives.

5 Inter stingly, concentrations of free radicals are maintained at high levels for more than ten minutes and tend to increase as tobacco smoke is aged. It is thus considered that these gas phase smoke oxidants are in a steady state as they are both continuously formed and destroyed. The latter reactions are similar to those noted to occur in smog, pointing to the extra noxious stimuli to  
10 primary and secondary smokers in polluted environments. In fact, environmental pollution in urban areas such as London, New York, Los Angeles and Mexico City promote gum disease and enhance the damage done to the oral cavity through the consumption of tobacco products. There is now also an established association between gum disease and coronary artery (heart)  
15 disease.

It is thus an object of the present invention to provide a gel, paste, gum or lozenge intended to be applied to the gums and teeth of a patient or taken orally as a means of preventing and ameliorating signs and symptoms and complications to the oral-pharyngeal cavity and mouth including the buccal  
20 mucosa and gums from damage caused by free radical species generating inflammation in gingival tissues.

These and further objects will be more readily apparent when considering the following disclosure and appended claims.

## 5 SUMMARY OF THE INVENTION

The present invention is directed to a gel, paste, gum or lozenge composition and a method of using it for preventing and reducing symptoms of periodontal disease in humans. The composition comprises a carrier and, as active ingredients, an effective amount of reduced glutathione and a source of  
10 selenium. Although the present composition is not considered a substitute for toothpaste, it can be applied to the gums in the case of a gel or paste by any well recognized means including the use of a tooth brush and other external applicators.

## DETAILED DESCRIPTION OF THE INVENTION

15 Human gums, are exposed to numerous environmental stimuli. Gums may be damaged by inhaled, ingested or chewed noxious substances, such as tobacco products, and may also be injured by systemic xenobiotics and by endogenous processes, such as inflammatory reactions. Innocent by-standers, the secondary smokers, may equally be adversely affected. Reactive oxidizing  
20 species, as induced by inhaled tobacco smoke, ozone and nitrous oxide, are important factors in generating free radicals and inducing inflammatory reactions resulting in the development of periodontal disease. The present invention has now recognized that such periodontal disease resulting from free radical generation can be in large prevented and its symptoms reduced by applying an  
25 effective amount of reduced glutathione and a source of selenium in a gel base or paste to the gum surface. The terms gel or paste include compositions wherein medicinal products may be included for specific oro-pharyngeal and odontologic conditions. Examples would include an anti-microbial agent such as those for treatment of oral candidiasis (thrush) or corticosteroid for management  
30 of oro-pharyngeal ulcers as both occur commonly in smokers and in patients with AIDS. In these oral condition, and particularly in patients with AIDS and their opportunistic infections, free radical species are indeed pathogenetic.

5 Thus, gels or paste with the compositions of this invention provide significant adjuncts of therapy in the management of oral conditions including gum disease.

It is now recognized that reducing the intracellular levels of GSH in cells increases their sensitivity to oxidant damage. Further, an L-cysteine delivery agent not only enhances endothelial cell GSH concentration, but also protects  
10 cells from damage from endogenous hydrogen peroxide. Without being bound to any particular theory, it is noted that reduced glutathione is employed in protecting cells against oxidative stress by itself being oxidized. Thus, L-glutathione must act in combination with other enzyme systems or cellular antioxidants in order to be reduced so that it may renew its role as a free radical  
15 scavenger. The enzyme glutathione reductase is vital to reduce the glutathione. GSH functions coordinately with the enzyme glutathione peroxidase which requires selenium as a cofactor to exert its biological antioxidant function. Selenium compounds, themselves, scavenge oxygen-centered radicals in vivo with reduced glutathione through glutathione peroxidase. By this physiologic  
20 method, selenium also functions as an anti-carcinogen. It is believed that selenium-GSH peroxidase catalyzes toxic hydrogen peroxide in the presence of reduced glutathione. This reaction reduces glutathione to oxidized glutathione (GSSG). In turn, the GSSG is reduced back to GSH by the enzyme GSH reductase thereby maintaining abundant cellular GSH to scavenge free  
25 radicals anew.

In summary, the major functions of reduced glutathione (GSH) in protection against lipid peroxidation involve three types of reactions, all inter-related and synergistic in combining non-enzymic scavengers and enzymic and dietary (vitamins C and E) provided antioxidants, to wit:

30 1. GSH with selenium co-factor glutathione peroxidase eliminate toxic peroxides.

2. GSH reduces oxidized forms of vitamin C which, in turn, maintains vitamin E in its reduced form promoting its metabolic functions. Thus, GSH



5 supports free radical reductions and free radical chain-terminating functions of the two nutrient antioxidants, vitamins C and E.

3. GSH functions through the glutathione S-transferases to detoxify reactive aldehydes created during the process of lipid peroxidation.

10 Glutathione and selenium act synergistically in vivo as they are both constituents of the same enzymatic system. GSH serves as a specific donor substrate while selenium, provided from alimentary sources or locally from topically administered preparations of selenium, selenoamino acids or selenium yeast extract, provides the prosthetic group for GSH peroxidase. The glutathione and selenium antioxidant functions are intrinsically related since, by  
15 keeping a peroxidase in action, the GSH and selenium contribute to the removal of the dismutation product of free oxygen radicals, namely, hydrogen peroxide. In a broad sense, GSH and selenium modulate free radical chains initiated or sustained by hydroperoxides. Selenium is used in the present invention for its role as an antioxidant as well as its anticarcinogenic and antimutagenic  
20 properties. Thus, selenium-glutathione complex may lower the level of potentially damaging peroxide radicals that are generated from various carcinogenic promoting chemicals, including tar phase and gas phase tobacco smoke inhaled by-products, particularly side stream smoke, that smoke which flows while the cigarette is being held in between puffs.

25 Glutathione peroxidase, a group of water soluble enzymes, also catalyze the destruction of both aqueous and membrane-bound hydroperoxides. In dietary selenium deficiency, these enzyme levels are markedly decreased resulting in severe free radical damage to the tissues so involved. The other related antioxidant systems cannot make up for depressed local activity of  
30 selenium and selenium dependent enzymes; thus, the importance of providing selenium in these intra-oral antioxidant preparations, as well as ascertaining adequate nutritional supplements.

5 L-ascorbic acid (vitamin C) or its derivatives can be employed in these compositions primarily for their antioxidant activities. Stabilized vitamin C is employed so that it does not lose its physiological reducing activities because of its high susceptibility to oxidation.

Other components were also investigated as being useful in practicing  
10 this invention, for example, the sulphur containing amino acid cysteine, which is one of the three amino acid constituents of the tripeptide antioxidant glutathione (GSH).

Vitamin A is an essential nutrient. Relative vitamin A deficiency may adversely affect the skin and mucous membranes, including the mucosa of the  
15 oral cavity and the gingiva. These alterations are reversible on repletion with vitamin A or one of its derivatives.

The present preparations may include flavorings. Flavors may be based on oils of spearmint and peppermint. Other flavoring materials may include menthol, clove, cinnamon, wintergreen, citrus fruits, eucalyptus, aniseed and  
20 others commercially available for these flavoring purposes.

Flavors may range in concentrations depending on the product from about 0.1% to about 6.0% by weight of the total composition.

The gels or pastes of this invention may include bicarbonates with thickening agents in a concentration from 0.5% to 5.0% by weight. State of  
25 the art thickeners with bicarbonate and zinc salts include, but are not limited to chicle, xanthan, arabic, karaya or tragacanth gums. Alginates, carrageenans and cellulose derivatives such as sodium carboxymethyl, methyl, or hydroxy ethyl compounds are appropriate for the intended preparations; surfactants and abrasives may also be included. Alcohols will otherwise be avoided for their  
30 known risk factor for oral cancers. In order to decrease dental cavities and add flavor, without using metabolizable sugars, sweetening agents as saccharin, sodium cyclamate, sorbitol, aspartame and others may be used in concentrations from 0.005% to 5.0% by weight of the total composition, albeit

5 the polyol xylitol, is preferred. Xylitol has been shown to prevent dental caries and decrease gum disease, in part by reducing the putative oral bacteria, especially *Streptococcus mutans*.

Gels and dentifrices may contain fluoride anticaries compounds. These fluoride compounds, such as salts of sodium, potassium, calcium, magnesium,  
10 stannous and others have been known to protect teeth from developing cavities. Hence, the various fluorinated water systems throughout the country. Fluorides may be present in various amounts in the gels, pastes, gums or lozenges ranging from 0.01% to 3.0% by weight, preferably from 0.05% to 2.0% by weight, most preferably from 0.1% to 1.2% by weight. These  
15 sources of stabilized fluoride are taught in U.S. Patent No. 5,372,802 by Barrows et al., issued December 13, 1994 and which is herein incorporated by reference. These compositions should release from 25 to 2000 PPM of the fluoride ion. The aforementioned patent taught a peroxygen compound, a fluoride and a zinc compound to inhibit the decomposition of the former. Thus,  
20 fluorides, optionally may be incorporated in the gels or pastes of these antioxidant preparations to prevent and repair free radical induced gingival diseases.

Other dental antiplaque oral compositions for gels and pastes have been patented with other compounds as optional ingredients. Gaffar in U.S. Patent  
25 No. 5,472,685 issued December 5, 1995, which is also incorporated herein by reference, teaches that the triclosan antimicrobial and antiplaque benefit is enhanced in the presence of a phenolic flavoring agent, such as menthol, eucalyptol or thymol. Antimicrobials may optionally be included in these gels and pastes to help decrease or eliminate the putative bacteria which contribute  
30 to the inflammatory reactions in gingivitis and periodontitis.

Lactitol may also be used as a substitute for sucrose in sugar-free compositions. Lactitol is a disaccharide sugar alcohol derived from lactose, highly water soluble and of low hygroscopicity, making it a suitable non-caloric

5 swe tener for use in solid dosage forms. A number of compounds may be added in order to enhance the aromas or tastes of these preparations. These substances may also impart fragrances to the aforementioned. Grapefruit and citrus aroma and flavors have been included in smoking tobacco articles. Methyl phenyl pentanol derivatives have been used to augment and enhance  
10 aromas, such as in perfumes and colognes. Schreck patented these derivatives for use in tobaccos and tobacco articles in U.S. Patent No. 4,458,699, issued July 10, 1994, which is herein incorporated by reference. Floral, green, weedy, fruity, minty, citrusy, oriental and green pepper-like aroma and taste nuances are well known to those skilled in the art of flavors and fragrances for such  
15 compositions as in oral sprays, mouthwashes, mouth rinses, gels, dentifrices and other medicinal, nutritional or breath freshener products.

As an optional embodiment, the compositions herein described may also contain zinc or zinc compounds. The state of the art of oral care and hygiene has long recognized the value of zinc to neutralize oral malodor and the value of  
20 zinc ions for their anti-plaque and anti-calculus properties. Mouth rinses, mouthwashes, gels and dentifrices will thus complement the properties of the xylitol sweetener in oral and dental preventative care.

Various patents have described different zinc compounds and other complexes in oral compositions. Domke and Bergman taught an aqueous zinc-  
25 polyamide complex as a solution for control of halitosis, dental care and to decrease the astringency and metallic taste of zinc in the mouth in U.S. Patent No. 5,587,147, issued December 24, 1996, which is herein incorporated by reference. This patent discloses previous documents dealing with zinc salts such as zinc chloride, zinc phenol, zinc sulfonate, zinc citrate and other zinc  
30 complexes, some of which purportedly also exhibit oral antimicrobial activities. The zinc ion concentrations in these compositions will be at least 0.1% to 3.0% by weight and these will preferably be in an alkaline pH to avoid

5 demineralization of tooth enamel at acid levels. In any event, these aqueous compositions will not have a pH below 6, and preferably about 7.

In order to reduce plaque formation, a number of optional ingredients may be added to the composition of the present invention. This is important for dental plaque is not only a source of dental cavities, but also a significant  
10 pathogenetic mechanism for gingivitis and periodontitis. The latter two clinical entities are the main targets of the dentifrice ingredients of the present invention. The oral bacteria and other pathogens associated with dental plaque not only create inflammation with an outpouring of white blood cells (leucocytes) which generate free radical species, but these micro-organisms also  
15 secrete enzymes and endotoxins which aggravate inflammation in the gums (gingivitis). This causes the gums to bleed and to lose their toughness that promotes its separation from the teeth, leading to infectious pockets (periodontitis). Thus, optional ingredients in these compositions include abrasives to help reduce plaque, as taught by Rice in U.S. Patent No.  
20 5,716,601 (Feb. 10, 1998) which is herein incorporated by reference.

The gels and pastes of this invention can also include one or more of the following abrasives at an inclusion range level of 5 to 95% by weight, based upon the weight of the entire composition such as hydrated silica, calcium carbonate, sodium bicarbonate, bicalcium phosphate, bentonite clay or kaolin  
25 clay. The preferred level of hydrated silica is 10-30%, the most preferred range is 15-22%.

Another method of application of the gel, paste, gum or lozenge of the present invention is to incorporate the various antioxidants, minerals and amino acids in liposomes or other state of the art encapsulating vehicles, akin to  
30 nanospheres, glycospheres and others as used in topical compositions. Liposomes are lecithin spheres that form an oil protective membrane around the putative active ingredients of the composition. These carriers also deliver the active ingredients locally for their preventive and therapeutic functions as well

5 as systemically through buccal mucosal absorption. Unger and co-workers, in U.S. Patent No. 5,580,575, issued December 3, 1996, which is herein incorporated by reference, have taught therapeutic drug delivery systems comprising gas-filled liposomes which encapsulate the active preparation. Earlier, Chakrabarti and associates, in U.S. Patent No. 5,380,531, issued  
 10 January 10, 1995, which is also herein incorporated by reference, disclosed preparations comprising a lipid and a modified peptide for incorporation into liposomes. In addition, Knight et al. (U.S. Pat. '388) has taught small particle aerosol liposomes and liposome combinations for medical delivery uses.

As an example of the present invention, a paste can be formulated having  
 15 the following ingredients.

	<u>INGREDIENTS</u>	<u>% BY WEIGHT</u>
	glycerol	32.0
	magnesium carbonate	3.5
20	sodium fluoride	0.10
	zinc acetate	0.50
	L-glutathione	0.1
	L-selenomethionine	0.05
	ascorbic acid	1.50
25	N-acetyl cysteine	0.10
	vitamin E	0.25
	benzalkonium chloride	0.10
	xylitol as sweetener	0.25
	polyvinyl pyrrolidone	7.5
30	coloring agent	0.20
	flavor, as peppermint	0.20

Ascorbic acid or ascorbyl palmitate may be incorporated in protective liposomes, as well as the glutathione and cysteine, N-acetyl cysteine and/or other amino acids employed in dentifrice formulations. As noted, whitening agents such as sodium monofluorophosphate and antimicrobials may also be  
 35 comprised in these dental formulations.

Further, the following gel is exemplary of the present invention.

	<u>INGREDIENTS</u>	<u>% BY WEIGHT</u>
	glycerin	42.0
40	poloxamer	18.0
	xylitol	3.5

5	ascorbic acid	2.0
	sodium lauryl sulfate	1.2
	natural peppermint oil	1.0
	alpha-tocopherol	0.75
	green tea	0.5
10	calcium lactate	0.25
	selenomethionine	0.20
	sodium fluoride	0.20
	L-glutathione	0.10
	coloring agent	0.10
15	deionized water	balance

In addition, the following composition is characteristic of a suitable lozenge.

X ACTIVE INGREDIENTS:

20	L-glutathione	50 mg.
	N-acetylcysteine	50 mg.
	selenium (selenomethionine)	70 mg.
	alpha lipoic acid	25 mg.
	vitamin A acetate	5000 I.U.
	ascorbic acid	180 mg.
25	X alpha tocopheryl acetate	60 I.U.
	copper lysinate	2 mg.
	manganese picolinate	15 mg.
	zinc picolinate	15 mg.

OPTIONAL ANTIOXIDANTS:

30	X grape seed extract	25 mg.
	green tea extract	50 mg.
	coenzyme Q10	50 mg.

SUGAR SUBSTITUTES:

xylitol or lactitol

5 W claim:

1. A gel or paste composition for oral application to human gums to prevent and reduce symptoms of gum disease, said composition comprising a gel or paste-based carrier and, as active ingredients, an effective amount of reduced glutathione and a source of selenium.

10 2. The gel or paste composition of claim 1 wherein said source of selenium comprises a member selected from the group consisting of elemental selenium, a selenoamino acid, a selenium yeast extract and a selenium chelate.

15 3. The gel or paste composition of claim 2 wherein said selenoamino acid comprises a member selected from the group consisting of selenomethionine and selenocystine.

4. The gel or paste composition of claim 1 further comprising vitamin C as ascorbic acid or as a derivative of ascorbic acid.

20 5. The gel or paste composition of claim 1 further comprising vitamin E as alpha tocopherol.

6. The gel or paste composition of claim 1 further comprising superoxide dismutase.

7. The gel or paste composition of claim 1 further comprising vitamin A.

25 8. The gel or paste composition of claim 1 further comprising at least one amino acid selected from the group consisting of cysteine, methionine, taurine and arginine or their derivatives.



5           9.     A method of preventing and reducing symptoms of periodontal disease in humans, said method comprising applying to human gums a gel or paste and, as active ingredients, an effective amount of reduced glutathione and a source of selenium.

10           10.    The method of claim 9 wherein said source of selenium comprises a member selected from the group consisting of elemental selenium, a selenoamino acid, a selenium yeast extract and a selenium chelate.

          11.    The method of claim 10 wherein said selenoamino acid comprises a member selected from the group consisting of selenomethionine and selenocysteine.

15           12.    The method of claim 9 further comprising vitamin C as ascorbic acid or as a derivative of ascorbic acid.

          13.    The method of claim 7 further comprising vitamin E as alpha tocopherol.

          14.    The method of claim 9 further comprising superoxide dismutase.

20           15.    The method of claim 9 further comprising vitamin A.

          16.    The method of claim 10 further comprising at least one amino acid selected from the group consisting of cysteines, methionine, taurine and arginine.

          17.    The gel or paste of claim 1, further comprising a flavorant.

25           18.    The gel or paste of claim 17 wherein said flavorant is xylitol.

          19.    The gel or paste of claim 1 wherein said active ingredients are encapsulated in a liposome.

5           20.    The gel or paste of claim 1 further comprising an abrasive.

          21.    The gel or paste of claim 20 wherein said abrasive is a member selected from the group consisting of hydrated silica, calcium carbonate, sodium bicarbonate, dicalcium phosphate, bentonite clay and kaolin clay.

          22.    The gel of claim 21 further comprising fluoride.

10           23.    A gum composition for oral application to human gums to prevent and reduce symptoms of gum disease, said composition comprising a gum-based carrier and, as active ingredients, an effective amount of reduce glutathione and a source of selenium.

          24.    A lozenge composition for oral application to human gums to  
15 prevent and reduce symptoms of gum disease, said composition comprising a lozenge-based carrier and, as active ingredients, an effective amount of reduce glutathione and a source of selenium.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/22507

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/48, 49, 52, 401, 440, 441, 450, 464, 702; 514/458, 474, 725, 900, 902, 904, 944

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

search terms: glutathione, selenium, antioxidant, periodontitis, gingivitis

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 5,667,791 A (HERSH et al) 16 September 1997, col. 2, lines 65-67; col. 3, line 1; col. 4, lines 17-19, 36-40; col. 8, lines 22, 57; col. 14, lines 54-56.	1-7 ----- 8-24
Y	US 5,378,461 A (NEIGUT) 03 January 1995, col. 9, lines 1-3; Abstract.	8-16
Y	US 5,032,384 A (YEH et al) 16 July 1991, col. 2, lines 31-38.	9-16
Y	US 5,709,873 A (BAR-SHALOM et al) 20 January 1998, col. 8, lines 18-23, 47-50, 59-67; col. 12, line 44.	17-24

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T"

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X"

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y"

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&amp;"

document member of the same patent family

Date of the actual completion of the international search

19 NOVEMBER 1999

Date of mailing of the international search report

23 DEC 1999

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MARINA LAMM

Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/22507

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

A61K 9/68, 7/16, 7/18, 7/00, 9/28, 9/127, 9/20, 33/04, 31/355, 31/34, 31/07

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/48, 49, 52, 401, 440, 441, 450, 464, 702, 514/458, 474, 725, 900, 902, 904, 944